



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A01N 43/54 // (A01N 43/54, 41:04, 37:10, 37:06, 37:04, 37:02)</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/40682</b> <b>(43) International Publication Date:</b> 6 November 1997 (06.11.97)
<b>(21) International Application Number:</b> PCT/GB97/01141 <b>(22) International Filing Date:</b> 25 April 1997 (25.04.97) <b>(30) Priority Data:</b> 9608771.3                      27 April 1996 (27.04.96)                      GB <b>(71) Applicant (for all designated States except US):</b> AGREVO UK LIMITED [GB/GB]; Hauxton, Cambridge CB2 5HU (GB). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> STOCK, David [GB/GB]; Chesterford Park, Saffron Walden, Essex CB10 1XL (GB). BRIGGS, Geoffrey, Gower [GB/GB]; Chesterford Park, Saffron Walden, Essex CB10 1XL (GB). SIMPSON, Donald, James [GB/GB]; Chesterford Park, Saffron Walden, Essex CB10 1XL (GB). <b>(74) Agent:</b> WALDMAN, Ralph, David; AgrEvo UK Limited, Patent Dept., Chesterford Park, Saffron Walden, Essex CB10 1XL (GB).		<b>(81) Designated States:</b> AU, BG, BR, CA, CN, CZ, HU, IL, JP, KR, MX, NZ, PL, RO, RU, TR, UA, US, VN, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.          Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> FUNGICIDE SALTS  <b>(57) Abstract</b>  Combining pyrimethanil with an organic acid having a volatility of less than 2 Pa at 20 °C results in a product which has valuable physical and biological properties.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Fungicide salts

This invention relates to compounds having fungicidal activity.

5

Pyrimethanil is a known fungicide, having the chemical name, 2-anilino-4,6-dimethylpyrimidine. However it has a relatively high vapour pressure which restricts its use. We have found that combining pyrimethanil with certain acids confers certain advantages to the compound.

10

According to the invention there is provided a product obtained by combining pyrimethanil with an organic acid having a volatility of less than 2 Pa at 20°C.

15

It is generally preferred that the acid is present in at least a stoichiometric amount and in this case a salt is formed between pyrimethanil and the acid. Excess acid may be an advantage, e.g. in a molar ratio of acid to pyrimethanil of up to 2:1.

20

As stated above one advantage of the products of the invention in particular is that they have a reduced vapour pressure compared with the free pyrimethanil, which increases the persistence of the compound on the crop to be protected from fungal attack. The reduced volatility also reduces levels of fungicide in the atmosphere.

25

In many cases the products have reduced phytotoxicity to certain plants.

30

In some cases the salts have increased activity compared with the free pyrimethanil. Another advantage is that the products have physical and chemical properties which often make them amenable to provide better formulations than the free pyrimethanil. For example, the product of pyrimethanil with oleic acid is liquid which provides formulation advantages compared with the free pyrimethanil which is a solid.

Suitable organic acids with which the pyrimethanil can be combined are fatty acids, especially long chain fatty acids such as oleic acid and palmitic acid. Other

suitable organic acids include saccharin, sulfonic acids, such as camphorsulfonic acid, salicylic acid and jasmonic acid.

The products are useful in combating diseases for which pyrimethanil may be used  
5 e.g. *Botrytis* spp., especially *B. cinerea*, *Venturia* spp, *Alternaria* spp., and *Monolinia fructigena*. However the salt may also extend the useful activity to diseases such as mildews and particularly cereal powdery mildew (*Erysiphe graminis*) and glume blotch (*Leptosphaeria nodorum*).

10 The invention is illustrated in the following Examples.

#### Example 1

A solution of pyrimethanil (1.0 g), toluene (50 ml) and oleic acid (1.42 g) was allowed to stand overnight at room temperature. The toluene was evaporated  
15 under reduced pressure to give pyrimethanil oleate, as an oil. (compound 1)  
nmr data:

CDCl<sub>3</sub>  $\delta$  scale

	0.9	( 3H , t , CH <sub>3</sub> )
20	1.25 - 1.42	( 20H , m , 10 x CH <sub>2</sub> )
	1.62 - 1.76	( 2H , m , CH <sub>2</sub> )
	1.95 - 2.1	( 4H , m , 2 x CH <sub>2</sub> )
	2.34 - 2.42	( 8H , m , 2xCH <sub>3</sub> , CH <sub>2</sub> )
	5.3 - 5.42	( 2H , m , CH = CH )
25	6.47	( 1H , s , pyrimidine CH )
	7.0	( 1H , t , ArH )
	7.32	( 2H , t , ArH )
	7.72	( 2H , d , ArH )
	8.67	( 1H , br s , NH )

30

#### Example 2

A solution of camphorsulfonic acid (1.25 g) in ethanol (10 ml) was added slowly to a solution of pyrimethanil (1 g) in toluene (20 ml) and the mixture allowed to stand for 30 minutes at room temperature. The mixture was evaporated under  
35 reduced pressure and the residue recrystallised from a mixture of diisopropyl ether

and ethyl acetate to give pyrimethanil camphorsulfonate, m.p. 166-7°C.  
(compound 2)

In a similar way there was obtained

- 5 a) pyrimethanil saccharinate, m.p. 164-5°C. (compound 3)
- b) pyrimethanil 7-trifluoromethylsaccharinate, m.p. 233-5°C. (compound 4)
- c) pyrimethanil 4,7-dimethoxysaccharinate, m.p. 187-8°C. (compound 5)
- d) pyrimethanil 4-chloro-7-methoxysaccharinate, m.p. 244-6°C.  
(compound 6)
- 10 e) pyrimethanil p-toluenesulfonate, m.p. 200-2°C. (compound 7)
- f) pyrimethanil 2H-1-benzopyran-3-carboxylate, m.p. 126-7°C. (compound 8)
- g) pyrimethanil phenoxyacetate, m.p. 76-8°C. (compound 9)
- h) pyrimethanil phenylphosphonate, m.p. 126-8°C. (compound 10)
- i) dipyrimethanil malonate, m.p. 126-8°C. (compound 11)
- 15 j) dipyrimethanil phthalate, m.p. 144-6°C. (compound 12)
- k) pyrimethanil hydrogen phthalate, m.p. 149-51°C. (compound 13)

### Example 3

This Example illustrates relative persistence of the products of the invention  
20 compared with the free anilinopyrimidine.

Droplets (5 x 4 µl) of toluene solutions of radiolabelled pyrimethanil (0.05% w/v)  
were applied to microscope cover slips (13 mm diameter), which were positioned  
in Petri dishes. To some of the samples were added various fatty acids in molar  
25 ratios of pyrimethanil to acid of 1:1 and 1:2. The Petri dishes were left in a  
controlled environment room (20°C, 16 hours daylight) and after two days, slips  
were removed to determine how much pyrimethanil remained. This was done by  
transferring the slips to scintillation vials, each containing 10 ml of a dioxane  
based scintillation cocktail and measuring the amount of radiation by liquid  
30 scintillation counting. The results are as follows:

Table 1: Surface recovery of pyrimethanil after 2 days

Compound	Surface recovery (%)
Pyrimethanil + oleic acid (1:1 molar)	59.5
Pyrimethanil + oleic acid(1:2 molar)	77.9
Pyrimethanil + lauric acid (1:1 molar)	66.9
Pyrimethanil + lauric acid (1:2 molar)	80.1
Pyrimethanil + myristic acid (1:1 molar)	71.9
Pyrimethanil + myristic acid (1:2 molar)	63.3
Pyrimethanil + palmitic acid (1:1 molar)	61.6
Pyrimethanil + palmitic acid (1:2 molar)	71.5
Pyrimethanil	3.1

In a similar manner the example was repeated by adding saccharin to the pyrimethanil in the amounts shown (% w/v of the toluene solutions). Surface recovery measurements were made after 2 and 8 days

The results are as follows:

10 Table 2: Surface recovery of pyrimethanil after 2 days

Compound	Surface recovery (%)
Pyrimethanil + saccharin (0.05%)	57.9
Pyrimethanil (0.05%) + saccharin (0.1%)	90.3
Pyrimethanil (0.05%) + saccharin (0.2%)	95.8
Pyrimethanil (0.05%)	2.0

Table 3: Surface recovery of pyrimethanil after 8 days

Compound	Surface recovery (%)
Pyrimethanil (0.05%) + saccharin (0.05%)	43.6
Pyrimethanil (0.05%) + saccharin (0.1%)	80.0
Pyrimethanil (0.05%) + saccharin (0.2%)	94.0
Pyrimethanil (0.05%)	1.1

It will be seen that the addition of the various acids increases the persistence of  
5 the pyrimethanil.

The compounds of Examples 1 and 2 also demonstrate greater levels of  
persistence than the free pyrimethanil.

Example 4

5% Wetttable powder formulations of compounds were diluted with water to the desired concentration and sprayed over wheat test plants. One day later the plants parts were inoculated with appropriate test pathogens and kept under controlled environment conditions suitable for maintaining plant growth and development of the disease. After an appropriate time, the degree of infection of the plant was visually estimated. Five replicates were used for each dose of test compound. The results are as follows. The rates of a.i. (active ingredient) in the tables are based on free pyrimethanil.

a) Botrytis cinerea (assessed 7 days after inoculation)

Compound No	% Control of disease at	
	5 g ai/hl	2 g ai/hl
5	94.4	85.4
6	94.4	88.7
7	91.0	83.1
8	93.2	84.2
9	90.4	75.2
10	94.4	87.6
12	91.0	77.5
pyrimethanil (5% WP)	85.4	75.2
SCALA	78.6	55.0

SCALA is the commercial 40% SC formulation of pyrimethanil



b) *Erysiphe graminis f. sp. tritici* (assessed 7 days after inoculation)

Compound No	% Control of disease at	
	100g ai/ha	25g ai/ha
4	39.7	15.5
8	75.9	27.6
12	51.7	3.4
pyrimethanil (5% WP)	27.6	3.4
SCALA	0	15.5

c) *Leptosphaeria nodorum* (assessed 21 days after inoculation)

5

Compound No	% Control of disease at	
	100g ai/ha	25g ai/ha
5	32.8	15.2
7	22.2	15.2
8	36.4	25.8
12	39.9	15.2
pyrimethanil (5% WP)	11.6	1.0
SCALA	15.2	11.6

Claims

1. A product obtained by combining pyrimethanil with an organic acid having a volatility of less than 2 Pa at 20°C.
- 5 2. A fungicidal composition which comprises a product as claimed in claim 1 in admixture with an agriculturally acceptable diluent or carrier.
3. A method of combating phytopathogenic fungi, at a locus infested or liable  
10 to be infested therewith, which comprises applying to the locus a compound of a product as claimed in claim 1.

## INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/GB 97/01141

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 6 A01N43/54 //(A01N43/54,41:04,37:10,37:06,37:04,37:02)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
 IPC 6 A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	AGRONOMIE, vol. 14, 1994, pages 541-554, XP002039434 P. LEROUX: "Influence du pH, d'acides aminés et de diversss substnces organiques sur la fongitoxicité du pyriméthanyl (....) vis-à-vis de certaines souches de Botrytis cinerea. " see page 544, column 1, paragraph 2 - page 550, column 1, paragraph 1	1-3
X	WO 92 19104 A (MYCOGEN CORP) 12 November 1992 see page 8, line 5 - page 10, line 15	1-3
A	EP 0 642 735 A (BASF AG) 15 March 1995 see page 3, line 56 - page 4, line 12 --- -/--	1-3



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&\* document member of the same patent family

Date of the actual completion of the international search

2 September 1997

Date of mailing of the international search report

24.09.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax (+31-70) 340-3016

Authorized officer

Klaver, J

# INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/GB 97/01141

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>EP 0 269 916 A (BAYER AG) 8 June 1988  see page 4, line 31 - page 5, line 12  -----</p>	1-3

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/GB 97/01141

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9219104 A	12-11-92	US 5366995 A	22-11-94
		AU 673496 B	14-11-96
		AU 1892892 A	21-12-92
		CN 1069391 A	03-03-93
		EP 0586477 A	16-03-94
		JP 6507399 T	25-08-94
-----			
EP 0642735 A	15-03-95	AT 153830 T	15-06-97
		CA 2131476 A	14-03-95
		CN 1111933 A	22-11-95
		DE 59402994 D	10-07-97
		ES 2102111 T	16-07-97
		HU 69037 A	28-08-95
		JP 7165512 A	27-06-95
		US 5508283 A	16-04-96
		US 5589479 A	31-12-96
		US 5591747 A	07-01-97
-----			
EP 0269916 A	08-06-88	DE 3639902 A	01-06-88
		AU 8161487 A	26-05-88
		JP 63154679 A	27-06-88
		US 4845098 A	04-07-89
-----			